NITRO | Additive Products



Benefits: Extracts from natural materials in the form of aqueous solution that is not salty, pH balance, plants can be absorbed through the roots and leaves. Suitable for the growing period of plants and restore the tree after harvest. It can be used with all types of plant. **Ingredient**: cow dung 35%, mixed vegetable juice 15%

Utilization Ratio	Recommendations use of each plant					
	Durian, Mangosteen and all fruit trees	Rice, Corn and Potato	Vegetables, Cactus and Ornamental plant			
20 Litre	50-100 cc	50 cc	30 cc			
Instruction			s radius and the entire plant evenly every 7-10 days. e soil surface for maximum efficiency.			



Additive Products



Benefits: Extracts from natural materials in the form of aqueous solution that is not salty, has a neutral pH, plants can be absorbed through the roots and leaves. Suitable for collecting food to prepare for flowering and the fruiting period until the harvest. It can be used with all types of plant.

Ingredient: Chicken dung and Guano 35%, mixed fruit juice 15%

Utilization Ratio	Recommendations use of each plant					
	Weather	Durian, Mangosteen, All fruit trees Rice, Corn and Potato	Longan	Salacca	Vegetable and Ornamental plant Weather	
	Normal Weather	30-50 cc	40 cc	60 cc	10 cc	
20 Litre	Heavy rain 40-60 cc 50 c	50 cc	70 cc	15 cc		
Instruction	Should be sprayed leaves and fruit continuously every 7-10 days by spraying into a small mist. canopies radius or soil surface.* Enough be wet over every part of the plant. Can be mixed with the sprinkled water system. *Spray the soil surface for maximum efficiency. *High humidity weather will use an additional 10-20% of the amount of the substance than usual.					



Additive Products



Benefits: Extracts from natural raw materials in the form of aqueous solution. Suitable for boosting immunity strengthen the roots, leaves, stems of plants **Ingredient**: Mangosteen extract 100%

	Introduction of use				
Suitable for	Utilization Ratio		Instruction		
	1 litre	20 litre	Histaction		
stems, branches and soil surface.		20-100 cc	1. Should be used consecutively at least 3-5 times. 2. Each time the period of use "TKO" no more than 1-7 days apart. 3. Can be used consecutively until the plant is strong. 4. Use TKO dosage increased or decreased can be considered as appropriate.		